

THE STRUCTURE OF ARRHYTHMIAS IN PATIENTS DURING RHINOLOGICAL SURGERY

I.K. Kalmykov¹, V.I. Popadyuk¹, P.D. Pryanikov², I.O. Efimenkov¹, N.D. Kuznetsov¹, A.O. Bykova¹

¹ Peoples' Friendship University of Russia (RUDN University), Moscow, Russia

² Russian Children's Clinical Hospital, Pirogov Russian National Research Medical University, Moscow, Russia
KIV: 0000-0001-6134-3080; ikastyro@gmail.com

Abstract: . In the present study, an attempt was made to study and evaluate the structure of cardiac arrhythmias in men and women using septoplasty under general anesthesia. Among all types of cardiac arrhythmias, premature atrial complexes and premature ventricular complexes extrasystoles (PAC and PVC) were most often found and were detected in all patients. Atrial tachycardias (AT) in men were significantly more likely to occur in the periods before surgery, during and after it, than in women. PACs at night was significantly more common in women. Ventricular arrhythmias (VA) were significantly more common in women than in men. Moreover, PVCs in women were detected significantly more often in all periods of observation. ATs and VAs in men were significantly more common after surgery. VAs in women were significantly more likely to occur before surgery than after it. Thus, the results of the study showed that of the considered cardiac arrhythmias during septoplasty under general anesthesia much more frequently in men and women at all stages of the survey found PVCs, and the women of their significantly higher than in men. Cardiac arrhythmias (PAC and PVC) predominated mainly in the postoperative period with the exception of PVCs in women, which was higher in the preoperative period. Reducing the number of arrhythmias observed during operation, possibly due to the impact of drugs for general anaesthesia.

Keywords: septoplasty, arrhythmia, stress, general anesthesia.

DOI: 10.25792/HN.2022.10.2.S1.47-49

For citations: Kalmykov I.K., Popadyuk V.I., Pryanikov P.D., Efimenkov I.O., Kuznetsov N.D., Bykova A.O. The structure of arrhythmias in patients during rhinological surgery. *Head and neck. Russian Journal.* 2022; 10 (2, Suppl. 1): 47-49

Introduction and aim. Septoplasty is a powerful surgical stressor [1-3]. After surgical interventions in the nasal cavity in animals, stress reactions occur due to an increase in predominantly nociceptive impulses that occur at the site of injury [4, 5]. As a rule, patients experience various types of cardiac arrhythmias and chronic hypoxia [6]. During surgical correction of deviated of nasal septum (septoplasty), which are a powerful stress factor, can cause arrhythmias [3, 5].

In the present study, an attempt was made to study and evaluate the structure of cardiac arrhythmias in men and women using septoplasty under general anesthesia.

Patients and Methods. For the period from June to October 2019, septoplasty was performed for 38 patients (20 men and 18 women, 18-55 years old) about the deviated of the nasal septum. Surgery was performed under general anesthesia using sol. Phentonyli (30 mkg / ml), sol. Midozolami (5 mg / ml), sol. Propofoli (150 mg), sol. Nimbexi (6 mg), sol. Traneksami (1000 mg), sol. Atropini (0.5 mg), sol. Ketonali (100 mg), sol. Cerucali (10 mg). Patients underwent holter ECG monitoring using a MT-101 Shiller (Switzerland) Halter, which began two hours before surgery and lasted for 24 hours. According to the results of the study, an analysis of the types of rhythm disturbances in each patient was carried out.

Results. Among all types of cardiac arrhythmias, premature atrial complexes and premature ventricular complexes extrasystoles (PAC and PVC) were most often found and were detected in all patients. Atrial tachycardias (AT) in men were significantly more likely to occur in the periods before surgery, during and after it, than in women. PACs at night was significantly more common in women. Ventricular arrhythmias (VA) were significantly more common in women than in men. Moreover, PVCs in women were detected significantly more often in all periods of observation. ATs and VAs in men were significantly more common after surgery. VAs in women were significantly more likely to occur before surgery than after it (tabl. 1).

Discussion. The nasal cavity contains the afferent section of many reflex arcs, which, passing through the autonomic nervous system, end in the heart [2-4]. Thus, manipulations in the nasal cavity can lead to dysregulation of cardiac activity and cause various kinds of arrhythmias [7, 8]. In experimental studies on the septoplasty simulation, it was shown that the application of trauma to the nasal septum mucous membrane entails an imbalance in the autonomic nervous system [9, 10], which is provoked by surgical damage in the maxillofacial region [10, 11]. The involvement of the central nervous system during stress reactions in the regulation of cardiac activity after septoplasty is also confirmed by morphological and physiological studies of the hippocampus formation [12-14]. To prevent the development of such complications of cardiac activity as arrhythmias, it is necessary to carefully assess the intraoperative risks, comorbidities of patients with

deviated nasal septum, especially from the cardiovascular system [2, 3, 15], and also clearly determine the presence of predictors of the development of postoperative pain syndrome [15].

Conclusion. Thus, the results of the study showed that of the considered cardiac arrhythmias during septoplasty under general anaesthesia much more frequently in men and women at all stages of the survey found PVCs, and the women of their significantly higher than in men. Cardiac arrhythmias (PAC and PVC) predominated mainly in the postoperative period with the exception of PVCs in women, which was higher in the preoperative period. Reducing the number of arrhythmias observed during operation, possibly due to the impact of drugs for general anaesthesia.

Conflict of interest. The authors declare no conflict of interest.

Tables 1. Type of arrhythmia in men and women during septoplasty. *- more than other sex ($p<0.01$); † - more than before septoplasty ($p<0.01$); ‡ - less than before septoplasty ($p<0.01$).

		men				women			
		before surgery	surgery	after surgery	night	before surgery	surgery	after surgery	night
Atrial tachycardias	PAC	8,25±2,4*	2.9±1.3‡	29.7±8.1†	11.6±4.1	4.6±1.4	3.1±1.4†	23.4±4.9†	19±5.6*†
	Couplet	0,4±0,13	2.1±1.2*†	3.6±1.7†	1.7±0.6†	1.4±0.6*	0.5±0.4†	2±0.6	3.4±2
	Triplet	0	0.1±0.1	2.9±1.7*†	0.6±0.2†	0	0.1±0.13	1.1±0.8†	2.9±2.4†
	SVTach	0,25±0,16*	0.3±0.2*	0.6±0.41	0.1±0.1	0	0	1.4±1†	0
	Bigem	0,06±0,05*	0	0.1±0.07	0.1	0	0.25±0.26	0.5±0.5	0
Ventricular arrhythmias	PVC	16.9±3,6	21.6±5.7	35.25±11†	18.75±7.8	90.4±10.6*	46.9±5.6*‡	72.5±12.4*†	38.9±10.9*‡
	Couplet	1.25±0,5	2.4±1.34	3.06±1.7†	0.9±0.6	8.6±3.5*	3±1.14‡	4.25±1.8‡	1.5±0.8‡
	Triplet	0.06±0,05	0.4±0.25†	0.56±0.3†	0.25±0.1†	2.4±0.9*	1±0.45*‡	0.75±0.5‡	0.25±0.26‡
	VTach	0.25±0.12	0.13±0.1†	0.19±0.15	0.06±0.05†	1.6±0.75*	0.5±0.28*‡	0.13±0.13‡	0‡
	Bigem	0.13±0.07	0.06±0.07	0.06±0.05†	0.06±0.05†	0.25±0.17	0.25±0.17*	0.5±0.3*	0.5±0.4

REFERENCES

1. Pustovit O.M., Nasedkin A.N., Egorov V.I., Isaev V.M., Isaev E.V., Morozov I.I. Using ultrasonic cavitation and photochromotherapy to increase nasal mucosa reparation process after septoplasty and submucous vasotomy of the inferior nasal turbinates. Golova I Sheya = Head and neck. Russian Journal. 2018;6(2):20–26 (in Russian).
 2. Kastyro I.V., Torshin V.I., Drozdova G.A., Popadyuk V.I. Acute pain intensity in men and women after septoplasty. Russian Open Medical Journal. 2017. 6 (3): 1-6.

3. Popadyuk V.I., Kastyro I.V., Ermakova N.V., Torshin V.I. Septoplasty and tonsillectomy: acute stress response as a measure of effectiveness of local anesthetics. Vestn Otorinolaringol. 2016; 81(3): 7-11.
 4. Kastyro I.V., Inozemtsev A.N., Shmaevsky P.E., Khamidullin G.V., Torshin V.I., Kovalenko A.N., Pryanikov P.D., Guseinov I.I. The impact of trauma of the mucous membrane of the nasal septum in rats on behavioral responses and changes in the balance of the autonomic nervous system (pilot study). J. Phys.: Conf. Ser. 2020; 1611 (012054)
 5. Kastyro I.V., Reshetov I.V., Khamidulin G.V., Shmaevsky P.E., Karpukhina O.V., Inozemtsev A.N., Torshin V.I., Ermakova N.V., Popadyuk V.I. The Effect of Surgical Trauma in the Nasal Cavity on the Behavior in the Open Field and the Autonomic Nervous System of Rats Doklady Biochemistry and Biophysics. 2020; 492: 121–123.
 6. Kastyro I.V., Reshetov I.V., Popadyuk V.I., Torshin V.I., Ermakova N.V., Karpukhina O.V., Inozemtsev A.N., Khamidulin G.V., Shmaevsky P.E., Sardarov

G.G., Gordeev D.V., Scopich A.A. Studying the physiological effects of a new model of septoplasty in rats. Head and neck. Russian Journal. 2020;8(2):33–38 (in Russian).
 7. Кастыро И.В., Гришина А.С. Изменения частоты сердечных сокращений на фоне различной анальгезии после септопластики. Российская оториноларингология. 2013; 1(62): 100-104.
 8. Kalmykov I.K., Torshin V.I., Ermakova N.V., Sinel'nikova A.N., Kastyro I.V. Otsenka ostrogo boleвого sindroma u patsientov posle septoplastiki pri primenenii razlichnykh taktik anestezii [Evaluation of acute pain

syndrome in patients after septoplasty and various anesthetic strategies]. *Ul'yanovskiy mediko-biologicheskii zhurnal*. 2021; 3: 97–110

9. Dragunova S.G., Reshetov I.V., Kosyreva T.F., Severin A.E., Khamidulin G.V., Shmaevsky P.E., A Inozemtsev.N., Popadyuk V.I., Kastyro I.V., Yudin D.K., Yunusov T.Yu., Kleyman V.K., Bagdasaryan V.V., Alieva S.I., Chudov R.V., Kuznetsov N.D., Pinigina I.V., Skopich A.A., Kostyaeva M.G. Comparison of the Effects of Septoplasty and Sinus Lifting Simulation in Rats on Changes in Heart Rate Variability. *Doklady Biochemistry and Biophysics*. 2021; 498: 165–169.

10. Kastyro I.V., Popadyuk V.I., Reshetov I.V., Kostyaeva M.G., Dragunova S. G., Kosyreva T.F., Khamidulin G.V., Shmaevsky P.E. Changes in the Time-Domain of Heart Rate Variability and Corticosterone after Surgical Trauma to the Nasal Septum in Rats. *Doklady Biochemistry and Biophysics*. 2021; 499: 247–250

11. Dolgalev Al.Al., Svyatoslavov D.S., Pout V.A., Reshetov I.V., Kastyro I.V. Effectiveness of the Sequential Use of Plastic and Titanium Implants for Experimental Replacement of the Mandibular Defect in Animals using Preliminary Digital Design. *Doklady Biochemistry and Biophysics*. 2021; 496: 36–39.

12. Kastyro I.V., Reshetov I.V., Khamidulin G.V., Shilin S.S., Torshin V.I., Kostyaeva M.G., Popadyuk . 2021; 10 (2): 34-41

V.I., Yunusov T.Y., Shmaevsky P.E., Shalamov K.P., Kupryakova A.D., Doroginskaya E.S., Sedelnikova A.D. Influence of Surgical Trauma in the Nasal Cavity on the Expression of p53 Protein in the Hippocampus of Rats. *Doklady Biochemistry and Biophysics*. 2021; 497: 99–103.

13. Torshin V.I., Kastyro I.V., Reshetov I.V., Kostyaeva M.G., Popadyuk V.I. The Relationship between P53-Positive Neurons and Dark Neurons in the Hippocampus of Rats after Surgical Interventions on the Nasal Septum. *Doklady Biochemistry and Biophysics*. 2022; 502: 30–35

14. Kostyaeva M.G., Kastyro I.V., Yunusov T.Yu., Kolomin T.A., Torshin V.I., Popadyuk V.I. Dragunova S.G., Shilin S.S., Kleiman V.K., Slominsky P.A., Teplov A.Y. Protein p53 expression and dark neurons in rats hippocampus after experimental septoplasty simulation. *Molekulyarnaya Genetika, Mikrobiologiya i Virusologiya (Molecular Genetics, Microbiology and Virology)*. 2022;40(1):39–45.

15. Кастыро И.В., Попадюк В.И., Торшин В.И. Острый болевой синдром после септопластики. М.: РУДН, 2021. 177с

16. Kastyro I.V., Romanko Yu.S., Muradov G.M., Popadyuk V.I., Kalmykov I.K., Kostyaeva M.G., Gushchina Yu.Sh., Dragunova S.G. Photobiomodulation of acute pain syndrome after septoplasty. *Biomedical Photonics*